

**Table 1.** Studies assessing the weight status of Cypriot adolescents (2000-2010)

Data Collection (year)	Study Sample	References	Total Sample Size (n)	RR (%)	Age (years)	Boys				Girls			
						n	%UW	%OW	%OB	n	%UW	%OW	%OB
2000	National <sup>1</sup>	Savva et al <sup>18</sup>	2,600 <sup>3</sup>	94.9%	10-18	835	-	21.3	7.1	859	-	16.5	4.5
2003	National <sup>1</sup>	Savva et al <sup>19</sup>	18,792 <sup>3,4</sup>	75.0%	11	774	-	21.4	8.8	751	-	21.8	7.0
2005	National <sup>1</sup>	Lazarou et al <sup>20</sup>	682 <sup>3</sup>	91.2%	10-14	336	-	19.0	11.6	346	-	18.5	8.4
2005	National	Christaki <sup>21</sup>	1,365	69.0%	11	466	3.1	19.1	16.9	476	5.7	15.9	11.5
2006	National	Lavithi <sup>22</sup>	1,500	56.9%	12-17	373	-	24.7	5.3	481	-	12.1	2.8
2007	National	Photiou <sup>23</sup>	4,271 <sup>3</sup>	NR	10-18	2,954	-	17.1	13.2	-	-	-	-
2007	Limassol	Loucaides & Jago <sup>24</sup>	247	95.5%	10-12	117	-	31.6	8.5	119	-	27.7	5.0
2008	National <sup>2</sup>	Loucaides et al <sup>25</sup>	1,966	91.6%	12-16	946	5.2	21.8	7.7	854	12.9	13.8	4.8

<sup>1</sup>Studies used in the sensitivity analyses (quality studies), <sup>2</sup>For the majority of the participants, weight and height were reported, not measured, <sup>3</sup>The total sample also included participants aged below 10 years old, <sup>4</sup>The study was cross-sectional at two time points and only the second sample, recruited after the year 2000, was used in the review. RR: Response Rate; NR: not reported; UW: underweight; OW: overweight; OB: obese.