

**Table 1.** Clinical characteristics of the subjects with (Group 1) and without (Group 2) autoimmune thyroiditis

	<b>Group 1 (n=19)</b>	<b>Group 2 (n=19)</b>	<b><i>p</i></b>
Age (years)	13.0 (10.1 – 14.7)	12.9 (10.5 – 14.9)	0.931
Gender (female)	12 (63%)	12 (63%)	0.999
Age at diagnosis (years)	8.1 (5.8 – 12.8)	9.0 (7.8 – 11.6)	0.665
Diabetes duration (years)	2.9 (0.4 – 5.3)	3.2 (1.5 – 5.1)	0.729
Weight SDS	-0.04 (-0.84 – 0.56)	-0.07 (-0.71 – 0.67)	0.840
Height SDS	-0.42 (-0.89 – 0.46)	-0.88 (-2.10 – -0.33)	0.146
BMI SDS	0.29 (-0.15 – 0.70)	0.25 (-0.58 – 1.32)	0.885
Exposure to smoke	9 (47%)	5 (26%)	0.138
Family history of autoimmune thyroiditis	5 (26%)	2 (11%)	0.209
Goiter	2 (10%)	0 (0%)	0.486
Pubertal subjects	13 (68%)	13 (68%)	0.999
Daily insulin dose (units/kg/d)	1 (0.7 – 1)	1.1 (0.8 – 1.6)	0.189

Results are given as median (25th - 75th percentiles) or n (%). SDS: standard deviation score; BMI: body mass index.