in thalassaemic patients.

Parameter Girls Boys p Total (n=19) (n=16) (n=35)

Table 2. Levels of IGF-I, IGFBP-3, ferritin and BMD measurements

	(n=19)	(n=16)	(n=35)
IGF-I SDS			
Mean±SD	$-0.35 \pm 0.95$	$0.14 \pm 0.82$	$-0.13 \pm 0.91$

 $-0.09\pm0.64$ 

 $1199 \pm 594$ 

242 - 2633

 $-0.41 \pm 0.95$ 

-1.8 - 0.9

-2.25 - 1.27 - 1.62 - 1.81

-0.95 - 1.35 - 0.95 - 1.17

 $-0.06 \pm 0.64$ 

 $1557 \pm 929$ 

627 - 4706

 $0.42 \pm 1.25$ 

-1.17 - 3.8

Range

Range

Range

SDS Mean±SD

Range

IGFBP-3 SDS Mean±SD

Serum ferritin (ng/L) Mean±SD

Lumbar BMD

0.109

0.877

0.177

0.034

-2.25 - 1.81

 $-0.08\pm0.63$ 

-0.95 - 1.35

 $1393 \pm 804$ 

242 - 4706

 $0.04 \pm 1.19$ 

-1.8 - 3.8